

Possible saturation effects at HERA and LHC in the k_T – factorization approach

A.V. Kotikov*, A.V. Lipatov[†] and N.P. Zotov[†]

**N.N. Bogoliubov Laboratory of Theor. Physics, Joint Institute for Nuclear Research,
141980 Dubna, Russia*

*[†]D.V. Skobeltsyn Institute of Nuclear Physics, M.V. Lomonosov Moscow State University,
119992 Moscow, Russia*

Abstract. We consider possible saturation effects in the structure function F_L at fixed W in HERA energy range and in bb –production at LHC energies in the framework of k_T –factorization approach.

Keywords: QCD, small x , BFKL evolution, k_T – factorization

PACS: 12.38.-t, 13.60.-r, 13.87.Ce

INTRODUCTION

Remarkable progress in recent years has resulted from the observation that the gluon density in a proton at small x grows as x decreases. Both DGLAP and BFKL evolution equations predict this rapid growth of the parton densities, thus demonstrating the triumph of perturbative QCD. It is, however, clear that this growth cannot continue for ever, because it would violate the unitarity constraint [1]. Consequently, the parton evolution dynamics must change at some point, and a new phenomenon must come into play. As the gluon density increases, non-linear parton interactions are expected to become more and more important, resulting eventually in the slowdown of the parton density growth (known under the name of "saturation effect") [1, 2]. The underlying physics can be described by the non-linear Balitsky-Kovchegov (BK) equation [3]. It is expected that these nonlinear interactions lead to an equilibrium-like system of partons with some definite value of the average transverse momentum k_T and the corresponding saturation scale $Q_s(x)$. This equilibrium-like system is the so called Color Glass Condensate (CGC) [4]. Since the saturation scale increases with decreasing of x , one may expect that the saturation effects will be more clear at the LHC energies. In the preasymptotic region ($k_T \geq Q_s(x)$) of energies the heavy quark production is described more adequately [5] by the so called k_T –factorization approach [1, 6]. At $k_T < Q_s(x)$ the k_T –factorization approach gives us a chance to account for saturation effects (if they are under control the BK equation) using scale properties of the dipole model, which is equivalent the k_T factorization [7].

Here we demonstrate a description of possible saturation effects in the framework of k_T –factorization approach. As example of that we consider the structure function F_L at fixed W and low Q^2 in HERA energy range and bb –quark production at LHC energies.




FIGURE 1. The Q^2 dependence of $F_L(x, Q^2)$ (at fixed $W = 276$ GeV). The experimental points are from [8]. Solid curve is the result of the k_T -factorization approach with the GLLM unintegrated gluon distribution from [9].

THEORETICAL FRAMEWORK AND NUMERICAL RESULTS

The SF F_L at fixed W in HERA range

The SF $F_L(x, Q^2)$ is driven at small x primarily by gluons and in the k_T -factorization approach it is related in the following way to the unintegrated distribution $\Phi_g(x, k_\perp^2)$:

$$F_L(x, Q^2) = \int_x^1 \frac{dz}{z} \int^{Q^2} dk_\perp^2 \sum_{i=u,d,s,c} e_i^2 \cdot \hat{C}_L^g(x/z, Q^2, m_i^2, k_\perp^2) \Phi_g(z, k_\perp^2), \quad (1)$$

where e_i^2 are charge squares of active quarks. The function $\hat{C}_L^g(x, Q^2, m_i^2, k_\perp^2)$ can be regarded as SF of the off-shell gluons with virtuality $k_\perp^2$ (hereafter we call them *hard structure function* by analogy with similar relations between cross-section and hard cross-section). They are described by the sum of the quark box (and crossed box) diagram contribution to the photon-gluon interaction (see, for example, Fig. 1 in [10]). To calculate the longitudinal SF $F_L(x, Q^2)$ we used the hard SF $\hat{C}_L^g(x, Q^2, m_i^2, k_\perp^2)$ obtained in Ref. [10] and different unintegrated gluon distributions.¹ Notice that the $k_\perp^2$ -integral in Eq. (1) can be divergent at lower limit, at least for some parameterizations of $\Phi_g(x, k_\perp^2)$. To overcome the problem we change the low Q^2 asymptotics of the QCD coupling constant within hard structure functions. We applied the so called "soft" version of "freezing" procedure [11], which contains the shift $Q^2 \rightarrow Q^2 + M^2$, where M is an additional

¹ Here we are interested in the parametrizations of unintegrated gluon functions which take into account saturations effects.




FIGURE 2. Azimuthal muon-muon correlations at LHC with different unintegrated gluon distributions

scale, which strongly modifies the infrared α_s properties. For massless produced quarks, ρ -meson mass m_ρ is usually taken as the M value, i.e. $M = m_\rho$. In the case of massive quarks with mass m_i , the $M = 2m_i$ value is usually used. We calculate the SF F_L as the sum of two types of contributions - the charm quark one F_L^c and the light quark one F_L^l :

$$F_L = F_L^l + F_L^c. \quad (2)$$

For the F_L^l part we used the massless limit of hard SF and restricted ourselves to the modification of the argument in the strong coupling constant of the hard SF only. We have shown [12] that our k_T -factorization results are in good agreement with the data for large and small part of the Q^2 range. However, there was some disagreement between the data and theoretical predictions at $Q^2 \sim 3 \text{ GeV}^2$. The disagreement comes from two possible reasons: additional higher-twist contributions, which are important at low Q^2 values², or/and NLO QCD corrections.

It was shown that the saturation (non-linear QCD) approaches contain information of all orders in $1/Q^2$, they resum higher-twist contributions [13]. The analysis of the behaviour of the longitudinal structure function $F_L(x, Q^2)$ in the saturation models was done in Ref. [14]³. In Fig.1 we demonstrate our k_T -factorization description of $F_L(Q^2)$ at fixed W with the unintegrated gluon distribution proposed in Ref. [9] which takes into account non-linear (saturation) effects. We see very well description of H1 experimental data at fixed W in all Q^2 region.

Saturation effects in $b\bar{b}$ -production at LHC

It was shown that the data on $b\bar{b}$ azimuthal correlations at Tevatron (measured as the decay muon ones, $d\sigma/d\Delta\phi^{\mu\mu}$) are much more informative to distinguish different unintegrated gluon distributions [15]. In Fig. 2 we show our predictions for the $b\bar{b}$ azimuthal

² Some part of higher-twist contributions was taken into account by the "freezing" procedure.

³ N.Z. thanks M.V.T. Machado for useful discussion of this problem.




FIGURE 3. The total transverse momentum distribution at LHC with different unintegrated gluon distributions

correlations at LHC obtained with different unintegrated gluon distributions (see, for example, [16]). The GBW and GLLM ones take into account saturation effects. We see that the latter two unintegrated gluon densities reduce to different behaviour of differential cross section $d\sigma/d\Delta\phi^{\mu\mu}$. Fig. 3 displays the total transverse momentum distribution, $d\sigma/dp_T^{bb}$, where $p_T^{bb} = p_T^b + p_T^{\bar{b}}$, at LHC energy obtained with the same unintegrated gluon densities. In this case the difference between the results with usual unintegrated gluon densities and ones obtained with account of the non-linear (saturation) effects is more dramatic.

CONCLUSIONS

We considered in the framework of k_T -factorization approach possible manifestation of the saturation effects at HERA and LHC. We shown that account of these effects (in particle, by the GLLM unintegrated gluon distribution) improves to a marked degree description of the F_L data at low Q^2 at HERA energy. We demonstrated also that the experimental data for $b\bar{b}$ azimuthal correlations and p_T^{bb} -distribution at LHC will give us additional possibilities to study non-linear (saturation) effects.

ACKNOWLEDGMENTS

N.Z. thanks DESY directorate for financial support and Organizing Committee for hostly and friendly atmosphere during the Symposium.

REFERENCES

1. L.V. Gribov, E.M. Levin and M.G. Ryskin, *Phys. Rep.*, **100**, 1-150 (1983).
2. A.H. Mueller and J. Qiu, *Nucl. Phys.*, **B 268**, 427-452 (1986).
L. McLerran and R. Venugopalan, *Phys. Rev.*, **D 49**, 2233-2241, 3352-3355 (1994); **D 50**, 2225-2233 (1994); **D 53**, 458-475 (1996); **D 59**, 094002-094021 (1999).
K. Golec-Biernat and M. Wusthoff, *Phys. Rev.*, **D 59**, 014017-014026 (1999); **D 60**, 114023-114041 (1999).
A.H. Mueller, hep-ph/9911289.
3. I.I. Balitsky, *Nucl. Phys.*, **B 463**, 99-160 (1996).
Y.V. Kovchegov, *Phys. Rev.* **D 60**, 034008-034015 (1999).
4. M. Gyulassy and L. McLerran, *Nucl. Phys.*, **A 750**, 30-63 (2005).
5. F. Gelis, R. Venugopalan, *Phys. Rev.*, **D 69**, 014019-014031 (2004).
D. Kharzeev, K. Tuchin, *Nucl. Phys.*, **A 735**, 248-266 (2004).
R. Baier, A.H. Mueller, D. Schiff, *Nucl. Phys.*, **A 741**, 358-380 (2004).
6. E.M. Levin, M.G. Ryskin, Yu.M. Shabelsky and A.G. Shuvaev, *Yad. Fiz.*, **53**, 1059-1076 (1991).
S. Catani, M. Ciafaloni and F. Hautmann, *Nucl. Phys.*, **B 366**, 135-188 (1991).
J.C. Collins and R.K. Ellis, *Nucl. Phys.*, **B 360**, 3-30 (1991).
7. V. Barone, M. Genovese, N.N. Nikolaev et al., *Phys. Lett.*, **B 326**, 161-167 (1994).
A. Bialas, H. Navelet and R. Peschanski, *Nucl. Phys.*, **B 593**, 438-450 (2001).
8. E.M. Lobodzinska, *Acta Phys. Polon.* **B 35**, 223-227 (2004); hep-ph/0311180.
9. G. Gotsman, E. Levin, M. Lublinsky and U. Maor, *Eur. Phys. J. C* **27**, 411-425 (2003).
10. A.V. Kotikov, A.V. Lipatov, G. Parente, N.P. Zotov, *Eur. Phys. J. C* **26**, 51-66 (2002).
11. N.N. Nikolaev and B.M. Zakharov, *Z. Phys.*, **C 49**, 607-618 (1991); **C 53**, 331-346 (1992).
12. A.V. Kotikov, A.V. Lipatov and N.P. Zotov, *JETP*, **101**, 811-816 (2005).
13. J. Bartels, K. Golec-Biernat, K. Peters, *Eur. Phys. J.*, **C 17**, 121-128 (2001).
E. Gotsman, E. Levin, U. Maor, L.D. McLerran, K. Tuchin, *Nucl. Phys.*, **A 683**, 383-405 (2001).
14. V.P. Goncalves, M.V.T. Machado, *Eur. Phys. J. C* **37**, 299-305 (2004).
15. S.P. Baranov, N.P. Zotov, A.V. Lipatov, *Phys. Atom. Nucl.* **67**, 837-845 (2004).
16. B. Andersson et al. (Small x Collaboration), *Eur. Phys. J. C* **25**, 77-101 (2002).